


TOO SICK FOR SCHOOL?



Below are some guidelines to help you make the decision about when to keep your child home from school. The recommendations are based on guidelines provided by the Center for Disease Control and local health professionals. They were developed to help prevent the spread of potentially contagious disease.




FEVER

With fever greater than 100.0° F. Student may return when fever-free for **24 hours** (WITHOUT use of fever-reducing medicine).



VOMITING/DIARRHEA

Any unexplained vomiting episode. May return **24** hours after last episode. Diarrhea = three or more unexplained episodes of watery or loose stool in **24** hours **OR** sudden onset of loose stools. May return 24 hours after last episode.



COUGH

Serious, sustained coughing, shortness of breath, or difficulty breathing.



RASH

Any new rash accompanied by a fever. May return after rash goes away or clearance given by a health care provider.



SKIN LESIONS/SORES

Drainage from a sore that cannot be contained within a bandage OR sores are increasing in size OR new sores are developing day-to-day.



OTHER

Symptoms that prevent the student from active participation in usual school activities OR student is requiring more care than school can safely provide.

Home is the best place for a child who is ill. If your child is sick with a diagnosed communicable disease, please notify the school as soon as possible. This notification will greatly assist others who, due to medical reasons and/or treatments, have weakened immune systems and may require immediate and specialized care.